

Nest Survey Report

Date: 06/16/2017

Biologist(s): Amy Trexler

Survey Area/Location/Structures:

Stations: 26 to 55

(Segment 5 - Sections 2, 3, and 4)

Stations 268 to 301

(Segment 5 - Sections 16, 17 and 18)

Proposed Construction within Survey Area:

Proposed construction activities include pot-holing and excavation and trenching for installation of new underground 230kV line and vaults.

Is Vegetation Clearing Required? Yes:	No: 🗵		
Was Complete Survey Coverage Achieved	? Yes: ⊠	No: 🗌	
Details of modifications to the survey area	if complete	coverage was not achieved:	N/A
Survey Conditions:	g, ,	.	
(T)*	Start	End	
Time	8:30 am	11:15 am	
Temperature (° F):	67 °	77 °	
Wind Speed (mph):	0-5 mph	0-5 mph	
Cloud Cover (%):	0%	0%	
Precipitation Type (if any) and duration:	n/a	n/a	

Habitat(s) and Vegetation Description within Survey Area:

- o Southern coast live oak riparian forest
- Southern willow riparian forest
- o Coastal sage scrub
- o Ornamental
- o Disturbed

Description of Survey Methodology:

The biologists conducted active and passive surveys within a survey buffer of 0.25 mile for white tailed kite, 500 feet for raptors, coastal California gnatcatcher, and least Bell's vireo, 250 feet for passerine birds in open space, and 150 feet for common passerines in residential, commercial and industrial areas. Active survey methods included walking meandering transects through the habitats while observing bird behavior with the aid of binoculars and directly searching in vegetation, the ground, the towers/poles, and other potential nest substrates. Passive survey methods included stationary observation periods from select vantage points that provided maximum visibility of the survey areas, using binoculars as necessary. If potential nesting behavior was observed within the survey buffers, specific shrubs were directly searched for nests in the areas where birds may have been observed exhibiting higher levels of activity or potential breeding behavior. All potential raptor nesting areas within the survey buffers were



searched directly and/or with the aid of binoculars. Visibility, access, time of year and weather conditions were all conducive to collecting comprehensive breeding data, and ample time was spent surveying all potential nest sites.

Suitable Raptor Nesting Habitat: Suitable CAGN Nesting Habitat: Suitable LBV Nesting Habitat: Survey Res	Yes: ⊠ Yes: ⊠ Yes: ⊠	No:
Nest(s) Located (complete table below if yes)*: Include previously located nests and current status)	Yes: 🖂	No: 🗌

Nest ID¹	Species ²	Listing Status ³	Nest Stage ⁴	Observation Notes ⁵	Latitude (decimal degrees)	Longitude (decimal degrees)
03212017_BSL_03	Red-tailed hawk (RTHA)	Common	Nestling/ Fledged	3/29- Adult observed on nest in western sycamore slightly above street level. Mate foraging nearby. 4/13- Adult observed sitting on nest. 5/12- Observed 2 chicks sitting in the nest. Chicks are approximately 14 days old. 5/23- Observed two chicks in the nest and the adult sitting in a nearby tree. 6/5- One chick sitting on a branch near the nest, close to fledging. The second chick has already fledged the nest. 6/13- One chick sitting on a branch near the nest, close to fledging. 6/16- Observed one chick laying in the nest. The other chick was in a nearby tree.	32.890656	-117.194337

^{1 –} Date (mmddyyy)_Biologist Initials (ABC)_Number ID (01) 2 – Include common name and four letter AOU species code



- 3 3 Federally Endangered (FE), Federally Threatened (FT), State Endangered (SE), State Threatened (ST), Species of Special Concern (SSC), Watch List (WL), Common
- 4-Building, Incubating, Nestling, Fledged, Complete/Inactive
- 5 Observation Notes: Item Carry (IC- nest material, food items, fecal sacs that indicate nesting in progress), Agitated/Territorial Behavior (ATB indicating potential nest sites or an intent to nest), Courtship Behavior (CB copulation, chasing flights, displays, etc.), Pair in Suitable Habitat (PSH utilizing all or portions of the buffer zone), Other

Avian Species Observed (complete common name):

Acorn woodpecker (ACWO), American crow (AMCR), Anna's hummingbird (ANHU), black phoebe (BLPH), bushtit (BUSH), California towhee (CALT), European starling (EUST), house finch (HOFI), house wren (HOWR), mourning dove (MODO), red-tailed hawk (RTHA), song sparrow (SOSP), spotted towhee (SPTO), and white-throated swift (WTSW).

Additional Species Observed:

*Belding's Orange-throated Whiptail (Aspidoscelis hyperythra beldingi) 32.89184, -117.19215

<u>Additional Notes (see Avian Species Observed and Observation Notes for definition of abbreviations):</u>

• Song sparrow and bushtit: pairs observed in suitable habitat and displaying territorial behavior

Nesting WTKI within 0.25 miles:	Yes:	No: 🖂
Nesting Raptor, CAGN, or LBV within 500 feet:	Yes:	No:
Nesting Passerine within 250 feet (open space only):	Yes:	No: 🛛
Nesting Passerine within 150 feet (residential,	Yes:	No: 🖂
commercial, residential areas):		

Establishment of Nest Buffer and Justification:

03212017_BSL_03- A nest buffer reduction request was approved by the CPUC on March 31, 2017. No changes or modifications to this nest buffer reduction request are required or recommended.